

Rec'd T/PTO 03 MAY 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number
WO 2004/041797 A1

(51) International Patent Classification⁷: C07D 277/82

Society, NSS Road, Gratkopar (West), Mumbai 400084,
Maharashtra (IN).

(21) International Application Number:

PCT/GB2003/004734

(74) Agents: WAIN, Christopher, Paul et al.; A.A. Thornton
& Co., 235 High Holborn, London WC1V 7LE (GB).

(22) International Filing Date:

3 November 2003 (03.11.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0225701.2

4 November 2002 (04.11.2002) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (*for all designated States except US*): CIPLA LTD [IN/IN]; 289 Bellasis Road, Mumbai Central, Mumbai 400 008 (IN).

(71) Applicant (*for MW only*): WAIN, Christopher, Paul [GB/GB]; A.A. Thornton & Co., 235 High Holborn, London WC1V 7LE (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): RAO, Dharmaraj, Ramachandra [IN/IN]; 4/403 Garden Enclave, Pokhran Road 2, Thane (West), Mumbai 400 601, Maharashtra (IN). KANKAN, Rajendra, Narayanrao [IN/IN]; A-3/5, NBD

(54) Title: PROCESS FOR PREPARING 2,6-DIAMINO-4,5,6,7-TETRAHYDRO-BENZOTHAZOLE

(57) Abstract: 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which is useful for making pramipexole, is made by: (i) reacting bromine with a solution of 4-acetamido-cyclohexanone in water to produce 2-bromo-4-acetamido-cyclohexanone; (ii) after step (i), adding thiourea to produce 6-acetyl amino- 2-amino-4,5,6,7-tetrahydro-benzthiazole; (iii) after step (ii), adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole dihydrobromide; and (iv) after step (iii), isolating 2,6-diamino-4,5,6,7 -tetrahydro-benzthiazole.

BEST AVAILABLE COPY

05/03/2005 TUE 07:31 [TX/RX NO 7138] 003

WO 2004/041797 A1

01533637

PCT/GB2003/004734

JC20 Rec'd PCT/PTO 03 MAY 2005

PROCESS FOR PREPARING 2,6-DIAMINO-4,5,6,7-TETRAHYDRO-BENZOTHAZOLE


This invention relates to a process for making 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, an intermediate useful in the production of pramipexole. The invention also relates to the synthesis of pramipexole.

(S)-4,5,6,7-tetrahydro-N6-propyl-2,6-benzothiazolodiamine (or (S)-2-amino-4,5,6,7-tetrahydro-6-(propylamino)benzothiazole), more commonly known as pramipexole, is used in both early and late Parkinson's disease as a dopamine agonist, to stimulate dopamine receptors in the brain. This has been described in EP 0 186 087.

EP 0 186 087 also describes the synthesis of various tetrahydro benzothiazoles, including pramipexole. In particular, the synthesis of pramipexole by the following reaction pathway is described. An initial reaction between bromine and 4-acetylamido-cyclohexanone is carried out in glacial acetic acid, with stirring for several hours, at room temperature. This is followed by the addition of thiourea under refluxing conditions. The reaction mixture is cooled, and crystals of 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole-hydrobromide are precipitated. The precipitate is filtered, then washed with water and acetone. The crystals are then dissolved in hydrobromic acid and the solution is refluxed for several hours. The solution is then concentrated by evaporation and the residue dissolved in methanol, from which crystals of 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide are formed. Subsequently, the 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide may be converted to pramipexole.

This synthesis is illustrated by the following reaction scheme:

2



This synthetic pathway involves separate reaction steps, each requiring different conditions, solvents, temperatures, etc. This necessitates a discontinuous process and more than one isolation step, which entails longer processing time, lower yields (product is lost during each isolation step), increased effluent load and increased solvent usage, in comparison with a continuous process.

We have now found a way of synthesising 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole from 4-acetamido-cyclohexanone, which avoids the multiple isolation steps used in the previously described processes.

According to one aspect of the present invention there is provided a method of synthesising 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which method comprises comprising: (i) reacting bromine with a solution of 4-acetamido-cyclohexanone in water to produce 2-bromo-4-acetamido-cyclohexanone; (ii) after step (i), adding thiourea to produce 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide; (iii) after step (ii), adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole; and (iv) after step (iii), isolating 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole free base.

It is an important feature of the present invention that step (iii) is carried out without any isolation of the 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole produced in step (ii). Thus the entire synthesis can be carried out in a single reaction vessel. Preferably at least three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.

Prior to step (i), the method may comprise the step of oxidising 4-acetamido-cyclohexanol to produce 4-acetamido-cyclohexanone. This step may be carried out in the same reaction vessel as subsequent steps (i) to (iv), thereby avoiding an additional isolation step.

The oxidation reaction may be carried out using oxidising agents including, for example, Jones reagent, sodium hypochlorite, manganese dioxide, pyridinium dichromate or potassium permanganate.

In step (i), the 4-acetamido-cyclohexanone solution and the bromine are preferably combined in the reaction vessel at a temperature in the range 5°C to 75°C, more preferably in the range 15°C to 40°C, and most preferably about room temperature (approximately 25°C). The bromine is preferably added dropwise to the 4-acetamido-cyclohexanone solution. After the bromine and the 4-acetamido-cyclohexanone solution have been combined, the mixture is preferably heated to a temperature in the range 30°C to 80°C, more preferably 40°C to 50°C,

WO 2004/041797

PCT/GB2003/004734

4

and most preferably about 45°C, and maintained at or near this temperature until the bromination is complete. The completion of bromination is indicated by the elimination of the characteristic brown colour of the bromine.

In step (ii), the temperature is preferably increased to 50°C to 95°C, more preferably to 70°C to 90°C, and most preferably to about 80°C.

In step (iii), the reaction mixture is preferably refluxed.

In step (iv), the reaction mixture is preferably cooled to 1°C to 35°C, more preferably to 5°C to 20°C, and most preferably to about 10°C, and the mixture is then neutralised. Typically the neutralisation is carried out with caustic lye solution (NaOH), although other alkalis may be used. Following neutralisation, the product, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, is isolated. The isolation may be carried out by filtration, centrifugation or any other suitable means. Following isolation, the product is preferably washed with chilled water.

The starting compound, 4-acetamido-cyclohexanone, may conveniently be formed by the oxidation of 4-acetamido-cyclohexanol.

The above described compound, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, is useful as an intermediate in the production of pramipexole and related compounds.

According to another aspect of the present invention there is provided a method of synthesising pramipexole, comprising the steps of: forming 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole by the method of the present invention, then converting it to pramipexole.

The conversion of 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole to pramipexole is well known in the prior art and is described, for example, in US 4,731,374. Any of the methods described in US 4,731,374, may be used in the present invention.

In one embodiment, the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole is converted to pramipexole by reaction with a propionyl halide, such as propionyl bromide, under suitable reaction conditions.

Both 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole and pramipexole have an asymmetric carbon atom, and exist as two distinct enantiomers: the S(-) isomer and the R(+)

isomer. The pharmacological activity of the S(-) isomer of pramipexole is, however, twice as high as that of the R(+) isomer, and the name pramipexole is commonly used to refer to the optically pure S(-) form. In this specification, "2,6-diamino-4,5,6,7-tetrahydro-benzthiazole" encompasses the R(+) and S(-) enantiomers individually and also encompasses any mixture thereof including the racemic mixture, and the term "pramipexole" encompasses the R(+) and S(-) enantiomers individually and also the racemic mixture.

The resolution of a racemic mixture of 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole can be carried out after step (iv) above. Methods of resolution are known in the art. Alternatively, pramipexole racemate can be produced prior to resolution, then the mixture resolved, if desired.

The resolution of pramipexole racemate is described by Schneider and Mierau (*J. Med. Chem.* 30, 494 (1987)). This method uses the di-amino derivative of (\pm)-4,5,6,7-tetrahydro-N6-propyl-2,6-benzothiazolediamine as a substrate and L(+) tartaric acid as a resolution agent. Following resolution, optically active pramipexole has been prepared by two-step propylation of the single enantiomer of the di-amino precursor comprising the steps of reaction with propionanhydride followed by a reduction of the propionyl intermediate.

The synthesis of the present invention avoids the need to isolate intermediate compounds, and thus the yield is higher and the processing time is lower. Furthermore, owing to the absence of an organic solvent (such as acetic acid) the costs are lower, and the reaction conditions are milder – the milder reaction conditions also have a positive impact on product purity.

The invention will now be further described with reference to the following Examples.

Example 1

Bromine (112g) was added dropwise to a solution of 4-acetamidocyclohexanone (100g) in 500ml water at room temperature. The mixture was warmed to approximately 45°C and maintained at this temperature until the bromine colour had been lost. To this,

thiourea (125g) was added, and the mixture was heated to approximately 80°C. To this, aqueous hydrobromic acid (100ml) was added, and the contents of the reaction vessel were refluxed. The contents were then cooled to approximately 10°C, and neutralized with caustic lye solution. The product, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, was isolated by filtration, and washed with chilled water and dried. The product was off-white in colour, and the yield was approximately 60g in weight.

Example 2

To a solution of 4-acetamido-cyclohexanol (100 g) in acetone (1L) was added Jones reagent (prepared from 68.5 g chromic oxide, 105 g sulphuric acid and 400 ml water) at 10-15°C. The excess of the reagent was quenched by addition of isopropanol (400 ml) and the solvent was removed under reduced pressure. Ethyl acetate (600 ml) was added, the contents stirred for 10 minutes and the lower aqueous portion drained off. Ethyl acetate was concentrated under reduced pressure and the residue dissolved in water (500 ml). Bromine (112 g) was added dropwise and the further reactions were carried out as described in Example 1.

Example 3

To a suspension of 4-acetamido-cyclohexanol (100 g) in water (300 ml) was added a solution of 10% sodium hypochlorite (500ml) and the contents stirred at room temperature for 12 hours. To this was added liquid bromine (112 g) and further reactions carried out as described in Example 1.

It will be appreciated that the invention described above may be modified.

CLAIMS:

- 1 A method of making 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which method comprises the steps in sequence of: (i) reacting bromine with a solution of 4-acetamido-cyclohexanone in water to produce 2-bromo-4-acetamido-cyclohexanone; (ii) adding thiourea to produce 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole; (iii) adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole; and (iv) isolating 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole.
- 2 A method according to claim 1 wherein step (iii) is carried out without isolating the 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole produced in step (ii).
- 3 A method according to claim 1 or 2, wherein any three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.
- 4 A method according to claim 1, 2 or 3, wherein steps (i) to (iv) are carried out in a single reaction vessel.
- 5 A method according to claim 1, 2, 3 or 4, further comprising, prior to step (i), the step of oxidising 4-acetamido-cyclohexanol to produce 4-acetamido-cyclohexanone.
- 6 A method according to claim 5, wherein the step of oxidising 4-acetamido-cyclohexanol to produce 4-acetamido-cyclohexanone and at least three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.
- 7 A method according to any preceding claim wherein in step (i) the solution of 4-acetamido-cyclohexanone in water and bromine are combined at a temperature of from 15°C to 40°C.

WO 2004/041797

PCT/GB2003/004734

8

- 8 A method according to any preceding claim wherein, after the bromine and the 4-acetamido-cyclohexanone solution have been combined, the mixture is heated to a temperature of from 40°C to 50°C, and maintained at or near this temperature until the bromination is complete.
- 9 A method according to any preceding claim wherein, in step (ii), the temperature is increased to 70°C to 90°C.
- 10 A method according to any preceding claim, wherein step (iii) is carried out under refluxing conditions.
- 11 A method according to any preceding claim wherein, after step (iii) but before step (iv), the reaction mixture is cooled to 5°C to 20°C, then neutralised.
- 12 A method according to any preceding claim, further comprising the step of resolving the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole isolated in step (iv) into its R(+) and S(-) enantiomers and recovering the R(+) and/or S(-) enantiomer.
- 13 A method of synthesising pramipexole, comprising the steps of : forming 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole by a method according to any preceding claim, and converting it to pramipexole.
- 14 A method according to claim 13, wherein 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole is converted to pramipexole by reaction with a propionyl halide.
- 15 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises the R(+) enantiomer.

- 16 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises the S(-) enantiomer.
- 17 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises a racemic mixture.
- 18 A method according to claim 14, further comprising the step of resolving the pramipexole into its R(+) and S(-) enantiomers and recovering the R(+) and/or S(-) enantiomer.
- 19 A method of synthesizing 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole substantially as herein described with reference to the Examples.

INTERNATIONAL SEARCH REPORT

Intern: Application No
PCT/GB 03/04734

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D277/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 186 087 A (DR. KARL THOMAE GMBH) 2 July 1986 (1986-07-02) cited in the application the whole document, particularly page 5478, right-hand column, first paragraph	1-19
Y	ALLINGER J ET AL: "The conformers of 2-bromocyclohexanone" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 2, 1958, pages 64-74, XP002253277 ISSN: 0040-4020 the whole document, particularly paragraph bridging pages 71 and 72	1-19
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *B* document member of the same patent family

Date of the actual completion of the international search

20 January 2004

Date of mailing of the international search report

28/01/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

INTERNATIONAL SEARCH REPORT

Interna Application No
PCT/GB 03/04734

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ALLINGER N L ET AL: "Conformational analysis. II. The 2-bromo-4-t-butylcyclohexanones" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC, US, vol. 80, no. 20, 20 October 1958 (1958-10-20), pages 5476-5480, XP002253278 ISSN: 0002-7863 the whole document, particularly page 5478, right-hand column, first paragraph -----	1-19

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No
PCT/GB 03/04734

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 186087	A	02-07-1986	
		DE 3447075 A1	03-07-1986
		DE 3508947 A1	18-09-1986
		AT 45735 T	15-09-1989
		AU 583874 B2	11-05-1989
		AU 5154485 A	17-07-1986
		BG 62023 B2	30-12-1998
		BR 1100678 A3	13-10-1999
		CA 1263653 A1	05-12-1989
		CS 9104099 A3	16-09-1992
		DD 242230 A5	21-01-1987
		DE 3572485 D1	28-09-1989
		DK 590285 A	23-06-1986
		EP 0186087 A1	02-07-1986
		ES 8702787 A1	01-04-1987
		ES 8707513 A1	16-10-1987
		ES 8707514 A1	16-10-1987
		ES 8707515 A1	16-10-1987
		FI 855102 A ,B,	23-06-1986
		GR 853126 A1	22-04-1986
		HK 78692 A	23-10-1992
		HU 39736 A2	29-10-1986
		IE 58863 B1	17-11-1993
		IL 77415 A	19-03-1990
		JP 1854941 C	07-07-1994
		JP 5072907 B	13-10-1993
		JP 61155377 A	15-07-1986
		KR 9309791 B1	11-10-1993
		LU 90208 A9	06-04-1998
		MX 9202792 A1	30-06-1992
		NO 855195 A ,B,	23-06-1986
		NZ 214661 A	26-04-1990
		PH 24533 A	03-08-1990
		PT 81735 A ,B	01-01-1986
		SG 82492 G	04-12-1992
		US 4843086 A	27-06-1989
		US 4886812 A	12-12-1989
		US 4731374 A	15-03-1988
		ZA 8509731 A	26-08-1987

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.